Nema Note 4



Nematode Sampling Problems

www.ncagr.com/agronomi/uyrnem.htm

Please consider the appropriate comment below concerning the soil sample(s) you submitted for nematode assay.

Note Comment

- **4-1** The sample submitted was too small to obtain a reliable assay. Therefore, interpret the results with caution. For a reliable assay, a sample at least one quart in size is necessary.
- 4-2 The sample was either too wet or too dry and may not have provided a reliable assay. Soil moisture levels similar to those for cultivation are optimum for nematode sampling. Placing the samples in plastic bags prior to shipment to the laboratory will prevent drying.
- 4-3 Very few nematodes were recovered from this sample. Since you did not indicate a recent soil fumigation, we suspect that the sample did not provide reliable assay material. Therefore, interpret the results cautiously, especially if a B recommendation was given. The population may have been underestimated.

NCDA&CS
AGRONOMIC DIVISION
NEMATODE ASSAY SECTION

PHYSICAL_ADDRESS 4300 REEDYC REEK ROAD RALEIGH NC 27607-6465

MAILING_ADDRESS 1040 MAIL SERVICE CENTER RALEIGH NC 27699-1040

> PHONE: 919-733-2655 FAX: 919-733-2837

> > Dr. Weimin Ye Nematologist

Dr. Colleen Hudak-Wise Division Director

Steve Troxler Agriculture Commissioner

For Additional Assistance

- Call your NCDA&CS regional agronomist or the Agronomic Division office in Raleigh (919-733-2655).
- ➤ Visit the NCDA&CS Agronomic Division Web site at www.ncagr.com/agronomi/.
- Visit your county Cooperative Extension office.
- Refer to the following online publications.
 - Sampling for plant-parasitic nematodes (NCDA&CS Agronomic Division, 2006)
 - --- www.ncagr.com/agronomi/pdffiles/samnemas.pdf
 - Taking soil samples for nematode assay (NCDA&CS Agronomic Division, 2006)
 - page 2 of www.ncagr.com/agronomi/pdffiles/isnemr.pdf
 - Taking field samples for diagnostic nematode assay (NCDA&CS Agronomic Division, 2006)
 - page 2 of www.ncagr.com/agronomi/pdffiles/isnempd.pdf